

INTERIM RECOVERY PLAN NO. 203

# **KUNDIP WATTLE**

## **(*ACACIA RHAMPHOPHYLLA*)**

### **INTERIM RECOVERY PLAN**

**2005-2010**

Renée Hartley<sup>1</sup> and Sarah Barrett<sup>2</sup>

<sup>1</sup> Technical Officer, CALM Albany Work Centre, 120 Albany Hwy, Albany 6330.

<sup>2</sup> Flora Conservation Officer, CALM Albany Work Centre, 120 Albany Hwy, Albany 6330



Photo: Bruce Maslin

August 2005

Department of Conservation and Land Management  
Albany Work Centre,  
South Coast Region,  
120 Albany Hwy, Albany WA 6331

## FOREWORD

Interim Recovery Plans (IRPs) are developed within the framework laid down in Department of Conservation and Land Management (CALM) Policy Statements Nos. 44 and 50.

IRPs outline the recovery actions that are required to urgently address those threatening processes most affecting the ongoing survival of threatened taxa or ecological communities, and begin the recovery process.

CALM is committed to ensuring that Threatened taxa are conserved through the preparation and implementation of Recovery Plans (RPs) or IRPs and by ensuring that conservation action commences as soon as possible.

This IRP will operate from September 2005 to August 2010 but will remain in force until withdrawn or replaced. It is intended that, if the taxon is still ranked Endangered, this IRP will be reviewed after five years and the need further recovery actions assessed.

This IRP was given regional approval on 26 October, 2005 and was approved by the Director of Nature Conservation on 26 October, 2005. The provision of funds identified in this IRP is dependent on budgetary and other constraints affecting CALM, as well as the need to address other priorities.

Information in this IRP was accurate at August 2005.

## ACKNOWLEDGMENTS

The following people have provided assistance and advice in the preparation of this Interim Recovery Plan:

Anne Cochrane	Manager, CALM Threatened Flora Seed Centre
Andrew Brown	Threatened Flora Coordinator, CALM Species and Communities Branch
Malcom Grant	Conservation Officer, CALM Ravensthorpe

Thanks also to staff of the W.A. Herbarium for providing access to Herbarium databases and specimen information, and CALM Wildlife Branch for their assistance.

## SUMMARY

<b>Scientific Name:</b>	<i>Acacia rhamphophylla</i>	<b>Common Name:</b>	Kundip Wattle
<b>Family:</b>	Mimosaceae	<b>Flowering Period:</b>	August to September
<b>CALM Regions:</b>	South Coast	<b>CALM District:</b>	Albany Work Centre
<b>Shire:</b>	Ravensthorpe	<b>Recovery Team:</b>	Albany District Threatened Flora Recovery Team

**Illustrations and/or further information:** Brown, A., Thomson-Dans, C. and Marchant, N. (Eds). (1998) *Western Australia's Threatened Flora*. Department of Conservation and Land Management, Western Australia; Western Australian Herbarium (1998) FloraBase - Information on the Western Australian Flora. Department of Conservation and Land Management, Western Australia. <http://www.calm.wa.gov.au/science/>.

**Current status:** *Acacia rhamphophylla* was declared as Rare Flora in 1996 under the Western Australian *Wildlife Conservation Act 1950* and is currently listed as Endangered under both State legislation and the Commonwealth *Environment and Biodiversity Protection Act 1999*. At the time of listing it met Endangered under World Conservation Union Red List Criterion D2 (IUCN 2001), primarily due to their being just a single known population and an unknown number of mature plants. However it strictly meets VU D2 as it is known that there are approximately 2000 plants over 5 hectares and little evidence of decline.

**Description:** This erect, woody-stemmed shrub, up to 70 cm high, has densely crowded greyish-green, spreading phyllodes (flattened leaf stalks that function as leaves) that are 11 to 17 mm long. Each phyllode is prominently grooved and round ended, with a short point below the tip. The stems of the plant appear black due to a covering of short hairs and black recurved, bristly stipules that are 5 mm long. The globular yellow flower heads are 2.5 to 3 mm and are on stalks up to 12 mm long. They are solitary and held in the axils of phyllodes at the ends of the branches. The hard, thin, brittle and blackish pods are 10 to 15 mm long and semi-circular in cross-section.

**Habitat requirements:** *Acacia rhamphophylla* is found in open shrub mallee on stony slopes in well drained sandy clay on or near contact between serpentine and banded iron formations. Plants are prominent in disturbed areas and along a drainage line.

**Habitat critical to the survival of the species, and important populations:** The habitat critical to the survival of *Acacia rhamphophylla* comprises the area of occupancy of the known population; similar habitat within 200 metres of the known population; remnant vegetation that may link future populations; and additional nearby occurrences of similar habitat that do not currently contain the species but may have done so in the past and may be suitable for translocations. Given that this taxon is listed as Endangered it is considered that all populations are important populations.

**Benefits to other species/ecological communities:** The Ravensthorpe Range occurs within one of the fifteen National Biodiversity Hotspots. The Ravensthorpe Range is habitat for a number of endemic species and threatened species, and some twenty Priority taxa. Recovery actions put in place for *Acacia rhamphophylla* will benefit these species and reciprocally, recovery actions put in place for these species will benefit *A. rhamphophylla*.

**International obligations:** This plan is fully consistent with the aims and recommendations of the Convention on Biological Diversity. *Acacia rhamphophylla* is not specifically listed under any international treaty and therefore this plan does not affect Australia's obligations under any other international agreements

**Role and interests of indigenous people:** According to the Department of Indigenous Affairs Aboriginal Heritage Sites Register, six registered sites occur in close proximity to the *Acacia rhamphophylla* populations. The Department has welcomed any future consultation that will seek input and involvement from Indigenous groups that have an active interest in the areas that are habitat for *A. rhamphophylla*.

**Affected interests:** This species is located on Crown land under mining tenements.

**Social and economic impacts:** The implementation of this recovery plan has the potential to have some social and economic impact as the population is located on mining tenements. However, recovery actions refer to continued negotiations between stakeholders with regard to these areas.

**Evaluation of the Plan's Performance:** The Department of Conservation and Land Management (CALM), in conjunction with the Albany District Threatened Flora Recovery Team (ADTFRT) will evaluate the performance of this IRP.

**Existing Recovery Actions:** The following recovery actions have been or are currently being implemented:

1. All land managers have been notified of the location and threatened status of the species.
2. Volunteers and staff from the CALM Albany Work Centre regularly monitor populations.

### **Objectives**

The objective of this Interim Recovery Plan is to abate identified threats and maintain or enhance *in situ* populations to ensure the long-term preservation of the species in the wild.

**Criteria for success:** The number of individuals within the population remains stable or increases over the five years of the plan.

**Criteria for failure:** The number of individuals within the population decreases over the five years of the plan.

### **Recovery actions**

1. Coordinate recovery actions.
2. Monitor populations.
3. Liaise with land managers.
4. Implement fire management.
5. Collect seed.
6. Obtain biological and ecological information.
7. Conduct further surveys.
8. Investigate the methodology for future translocation(s).
9. Map habitat critical to the survival of the species.
10. Promote awareness.
11. Review the IRP and assess the need for further recovery actions.

## 1. BACKGROUND

### History

Kundip wattle was discovered in a single area during a flora survey in 1992 and, despite extensive surveys of similar habitat by the Ravensthorpe Wildflower Society and CALM staff, is still known only from the type population in the Ravensthorpe Range. The population is located on mining tenements, and mining and track maintenance pose the greatest threat. The current leaseholder is aware of the population and has assisted with conservation efforts to date.

### Description

This erect, woody-stemmed shrub, up to 70 cm high, has densely crowded greyish-green, spreading phyllodes (flattened leaf stalks that function as leaves) that are 11 to 17 mm long. Each phyllode is prominently grooved and round ended, with a short point below the tip. The stems of the plant appear black due to a covering of short hairs and black recurved, bristly stipules that are 5 mm long. The globular yellow flower heads are 2.5 to 3 mm and are on stalks up to 12 mm long. They are solitary and held in the axils of phyllodes at the ends of the branches. The hard, thin, brittle and blackish pods are 10 to 15 mm long and semi-circular in cross-section.

*Acacia rhamphophylla* appears most closely related to *A. laricina* and *A. cedroides* (Maslin 2001).

### Distribution and habitat

*Acacia rhamphophylla* is known from a single population, occupying approximately five hectares in the Ravensthorpe Range. It occurs in open shrub mallee on stony slopes in well drained sandy clay, on or near contact between serpentine and banded iron formations. While plants are most common in disturbed areas, they also occur in lower numbers under mature vegetation. Most plants are concentrated along a drainage line but some are also found upslope.

Associated species include *Eucalyptus cernua*, *E. pleurocarpa*, *E. transcontinentalis*, *Alyogyne hakeifolia*, *Beaufortia schaueri*, *Acacia durabilis*, *A. pinguiculosa*, *Cooperookia polygalacea*, *Hybanthus floribundus* and *Melaleuca* species.

The species was not found during detailed surveys of Bandalup Hill, approximately sixteen kilometres east of the known population. Similarly, no *Acacia rhamphophylla* juveniles have been found in neighbouring vegetation burnt during a 2000 wildfire.

### Biology and ecology

Little is known about the biology and ecology of *Acacia rhamphophylla*. Mature plants flower from August to September, though the juvenile period is unknown. In some species of *Acacia*, all of the flowers are hermaphrodite, while in others a percentage of the flowers on an individual are purely male. Unlike many other flower structures, acacia flowers have no complex morphological traits to exclude specific visitor taxa. As a result, acacias are visited by a large variety of insect and some bird pollinators and are vulnerable to exploitation by non-pollinators. Pollinators may favour certain *Acacia* species however, depending on such factors as number and density of flower heads. Pollinator limitation is a concern for endangered *Acacia* species in Australia (Stone *et al.* 2003).

Prior to *Acacia rhamphophylla* being recognised as Declared Rare Flora, mining activities resulted in some disturbance from fire and vehicle movements (<sup>3</sup>M. Grant, personal communication). The species appeared to recover well with regeneration of seedlings in the disturbance areas. This, coupled with its prolific flowering ability, suggests the species is capable of producing good quantities of viable seed (M. Grant, personal communication).

---

<sup>3</sup> Malcom Grant

Conservation Officer, CALM Ravensthorpe

Legumes are described as disturbance opportunists (Schwarten 1995). Fire is the most common environmental cue for breaking seed dormancy in the majority of *Acacia* species (Yates and Broadhurst 2002). In recent years, a significant number of deaths amongst mature *A. rhamphophylla* have been observed in the population. This may suggest a senescent population and reinforce the necessity for germination stimulants.

Research has shown that the majority of *Acacia* species are resistant to *P. cinnamomi*, however *A. rhamphophylla* has not been tested to date, nor have the two species (*A. laricina* and *A. cedroides*) thought to be most closely related to it (<sup>4</sup>B. Shearer, personal communication).

## Threats

*Acacia rhamphophylla* was declared as Rare Flora in 1996 under the Western Australian *Wildlife Conservation Act 1950* and is currently listed as Endangered under both State legislation and the Commonwealth *Environment and Biodiversity Protection Act 1999*. At the time of listing it met Endangered under World Conservation Union Red List Criterion D2 (IUCN 2001), primarily due to their being just a single known population and an unknown number of mature plants. However it strictly meets VU D2 as it is known that there are approximately 2000 plants over 5 hectares and little evidence of decline.

All areas occupied by *Acacia rhamphophylla* are affected or potentially affected by one or more threats identified in this IRP. Threats include:

- **Mining:** Population 1 occurs on Crown land subject to a live mining lease, a pending mining lease and a pending exploration licence. Future impacts of mining may include vegetation clearing, ground compaction, dust, introduction of weeds and pathogens such as *Phytophthora cinnamomi*, increased risk of fires and discharge of waste products and hazardous materials.
- **Inappropriate fire regime:** Poorly timed, intense and too frequent fire may be detrimental, as plants need to reach reproductive maturity to build up a seed bank. An estimation of the minimum desirable fire interval may be determined by doubling the primary juvenile period (time to first flower from germination, in 50% of the population) (Gill and Nichols 1989), however for *Acacia rhamphophylla*, this period is unknown. Equally, if the fire interval exceeds the longevity of the plants and the seed bank, population decline and extinction can occur (Yates and Broadhurst 2002).
- **Small population size:** As population size decreases, the population may become more vulnerable to extinction for three main reasons. Firstly, loss of genetic variation and increased inbreeding are considered to be associated with a reduction in the ability of a population to adapt to short-term environmental change. Secondly, small populations are more susceptible to chance events associated with demographic and environmental stochasticity. Finally, Allee effects may occur, whereby at some density or population size, reproductive capacity drops below a threshold and the organism can no longer replace itself (Hobbs and Yates 2003).
- **Climate change:** Long-term climate change may affect the *Acacia rhamphophylla* population given the predicted decrease in rainfall and increases in temperature and evaporation. It has been considered that those groups likely to be most affected by climate change include geographically localised taxon such as *A. rhamphophylla*, peripheral or disjunct populations, specialised species, poor dispersers, genetically impoverished species, and coastal communities (Peters & Darling 1985). Studies show that a decrease in rainfall and a shorter wet season may have been responsible for the reduced flowering, fruiting and seed production in a number of *Acacia* species over recent years (Yates and Broadhurst 2002)

## Summary of population land vesting, purpose and tenure

Population	Vesting	Purpose	Tenure
------------	---------	---------	--------

<sup>4</sup> Bryan Shearer

Principle Research Scientist, CALM Science

1. Mt Iron	Unvested	Common	Crown
------------	----------	--------	-------

### Summary of population information and threats

Pop. No. & Location	Year/No. plants	Condition	Threats
1. Mt Iron	1996 1000+	Healthy	Mining
	1997 2000+	Healthy	Track maintenance
	1999 2000+	Healthy	Fire
	2002 1000-2000	Healthy (few deaths)	
	2004 2000+/-	Healthy (<5% death)	

### Habitat critical to the survival of the species, and important populations

Given that this species is listed as Endangered under the Commonwealth EPBC Act, it is considered that all known habitat is habitat critical to the survival of the species. In addition all populations, including any translocated populations, are considered important to the survival of the species. Recovery actions include survey. Habitat is defined as the biophysical medium or media occupied (continuously, periodically or occasionally) by an organism or group of organisms or once occupied (continuously, periodically or occasionally) by an organism, or group of organisms, and into which organisms of that kind have the potential to be reintroduced (*Environment Protection and Biodiversity Conservation Act 1999*). The area of occupancy of the currently known *Acacia rhamphophylla* population has been mapped. However, other parts of the habitat critical to the survival of *A. rhamphophylla* have not been mapped and an action outlined in this Interim Recovery Plan is to map all habitat as defined above.

The habitat critical to the survival of *Acacia rhamphophylla* therefore comprises:

- the area of occupancy of known populations;
- areas of similar habitat within 200 metres of known populations that provide potential habitat for natural recruitment;
- remnant vegetation that surrounds and links populations (this is necessary to allow pollinators to move between populations) and
- additional occurrences of similar habitat that do not currently contain the species but may have done so in the past (these represent possible translocation sites).

### Benefits to other species/ecological communities

The Ravensthorpe Range is an area of high conservation value and occurs within one of the fifteen National Biodiversity Hotspots, which are areas of species richness and endemism, and areas under major threat (CALM 2004). The Ravensthorpe Range is habitat for a number of endemic species and Threatened species, including *Daviesia megacalyx* (En), *Marianthus villosus* (Vu) and some twenty Priority taxa, such as *Melaleuca stramentosa* (P1), *Pultenea* sp. Kundip (P1), *Melaleuca* sp. Kundip (P1), *Acacia larinia* var. *crassifolia* (P2), *Spyridium glaucum* (P3) and *Siegfriedia darwinioides* (P4). Recovery actions put in place for *Acacia rhamphophylla* will benefit these species and reciprocally, recovery actions put in place for these species will benefit *A. rhamphophylla*.

### International Obligations

This plan is fully consistent with the aims and recommendations of the Convention on Biological Diversity, ratified by Australia in June 1993, and will assist in implementing Australia's responsibilities under that Convention. However, as *Acacia rhamphophylla* is not listed under any international agreement, the implementation of other international environmental responsibilities is not affected by this plan.

### Role and interests of indigenous people

According to the Department of Indigenous Affairs Aboriginal Heritage Sites Register, the registered sites Claytup Surface Scatter, Kundip, Coujinup Surface Scatter, Gnamma Hole and North Jerdacuttup River 1



& 2 occur in close proximity to *Acacia rhamphophylla*. The Department has welcomed any future consultation that will seek input and involvement from Indigenous groups that have an active interest in the areas that are habitat for *Acacia rhamphophylla*, and this is discussed in the recovery actions.

### **Affected Interests**

The population occurs on Crown land under mining tenements.

### **Social and economic impacts**

The implementation of this recovery plan has the potential to have some minimal social and economic impact as the population is located on mining tenements. However, recovery actions refer to continued negotiations between stakeholders with regard to these areas.

### **Guide for decision-makers**

Section 1 provides details of current and possible future threats. Developments in the immediate vicinity of the population or within the defined critical habitat of *Acacia rhamphophylla* require assessment for the potential for a significant level of impact. No developments should be approved unless the proponents can demonstrate that they will not have a detrimental impact on the species, or its habitat or potential habitat, or the local surface and ground water hydrology.

### **Evaluation of the Plan's Performance**

The Department of CALM, in conjunction with the Albany District Threatened Flora Recovery Team will evaluate the performance of this recovery plan. In addition to annual reporting on progress against the criteria for success and failure, the plan is to be reviewed within five years of its implementation. Any changes to management and/or recovery actions made in response to monitoring results will be documented accordingly.

## **2. RECOVERY OBJECTIVE AND CRITERIA**

### **Objectives**

The objective of this Interim Recovery Plan is to abate identified threats and maintain or enhance *in situ* populations to ensure the long-term preservation of the species in the wild.

**Criteria for success:** The number of individuals within the population remains stable or increases over the five years of the plan.

**Criteria for failure:** The number of individuals within the population decreases over the five years of the plan.

## **3. RECOVERY ACTIONS**

### **Existing or completed recovery actions**

The land managers have been notified of the location and threatened status of *Acacia rhamphophylla*. The notification details the Declared Rare status of the species and the legal responsibility to protect it.

Staff at the CALM Albany Work Centre regularly monitor the population and liaise with the land holder.

### **Future recovery actions**



Where populations occur on lands other than those managed by CALM, permission has been or will be sought from appropriate land managers prior to recovery actions being undertaken. The following recovery actions are roughly in order of descending priority; however this should not constrain addressing any of the priorities if funding is available and other opportunities arise.

### 1. Coordinate recovery actions

The Albany District Threatened Flora Recovery Team (ADTFRT) is coordinating recovery actions for *Acacia rhamphophylla* and will include information on progress in their annual report to CALM's Corporate Executive and funding bodies.

**Action:** Coordinate recovery actions  
**Responsibility:** CALM (Albany Work Centre) through the ADTFRT  
**Cost:** \$3,000 per year

### 2. Monitor populations

Continue regular monitoring of *Acacia rhamphophylla*.

**Action:** Monitor populations  
**Responsibility:** CALM (Albany Work Centre)  
**Cost:** \$730 per year

### 3. Liaise with land managers

Staff from CALM Albany District will continue to liaise with current and future mining leasees to ensure populations on mining tenements are not accidentally damaged or destroyed and that the impacts of identified threats are minimised. Input and involvement will also be sought from Indigenous groups that have an active interest in areas that are habitat for *Acacia rhamphophylla*.

**Action:** Liaise with land managers  
**Responsibility:** CALM (Science Division and Albany Work Centre)  
**Cost:** \$600 per year

### 4. Implement fire management

A fire management strategy will be developed and implemented. The use of fire to stimulate recruitment will be considered if the population shows signs of decline.

**Action:** Implement fire management  
**Responsibility:** CALM (Albany Work Centre)  
**Cost:** \$2,600 in the first year

### 5. Collect seed

There is currently no *Acacia rhamphophylla* seed in storage, however preservation of germplasm is essential to guard against the possible extinction of wild populations. Seed is required to propagate plants for future translocations. Seed collection will be ongoing to obtain seed from as wide a range of individuals as possible to maximise the genetic diversity of *ex situ* material.

**Action:** Ongoing seed collection  
**Responsibility:** CALM (Albany Work Centre)  
**Cost:** \$1,530 per year

### 6. Obtain biological and ecological information

Improved knowledge of the biology and ecology of *Acacia rhamphophylla* will provide a better scientific basis for management of the wild populations. An understanding of the following is particularly necessary for effective management:

1. Disease susceptibility.
2. Soil seed bank dynamics and the role of various disturbances, competition and rainfall in germination and recruitment.
3. The pollination biology, phenology and seasonal growth of the species.
4. The population genetic structure, levels of genetic diversity and minimum viable population size.

**Action:** Obtain biological and ecological information  
**Responsibility:** CALM (Science Division and Albany Work Centre) through the ADTFRT  
**Cost:** \$24,000 per year for the final three years

## 7. Conduct further surveys

Surveys supervised by CALM staff, with assistance from local naturalists and wildflower society members, are to be conducted during the species flowering period (August to September). Similar habitat has not been extensively surveyed to date. Information on soil and vegetation types will be used to identify similar habitat to target for further survey.

**Action:** Conduct further surveys  
**Responsibility:** CALM (Albany Work Centre)  
**Cost:** \$2,500 per year

## 8. Investigate the methodology for future translocation(s)

Within the 5-year time frame of the plan, the best methodology for future translocations will be investigated. The most appropriate translocation site and procedure should be determined.

**Action:** Investigate the methodology for future translocation(s)  
**Responsibility:** CALM (Science Division and Albany Work Centre)  
**Cost:** \$2,500 per year

## 9. Map habitat critical to the survival of the species

It is a requirement of the EPBC Act (Section 207A) that spatial data relating to critical habitat be determined. Although habitat that is critical to the survival of the species is alluded to in Section 1, all the areas described have not yet been accurately mapped and will be addressed under this action. If additional populations are located, habitat critical to their survival will also be determined and mapped.

**Action:** Map habitat critical to the survival of the species  
**Responsibility:** CALM (Albany Work Centre)  
**Cost:** \$400 in first year

## 10. Promote awareness

The importance of biodiversity conservation and the need for the long-term protection of wild populations of this species will be promoted to the community through poster displays and the local print and electronic media. Formal links with local naturalist groups and interested individuals will also be encouraged.

**Action:** Promote awareness  
**Responsibility:** CALM (Albany Work Centre) through the ADTFRT  
**Cost:** \$900 per year

## 11. Review the IRP and assess the need for further recovery actions

If *Acacia rhamphophylla* is still ranked as Endangered at the end of the fourth year of the five-year term of this IRP, the plan will be reviewed and the need for further recovery actions assessed.

<b>Action:</b>	Review the IRP and assess the need for further recovery actions
<b>Responsibility:</b>	CALM (Species and Communities Branch and Albany Work Centre) through the ADFRT
<b>Cost:</b>	\$4,000 in the fifth year (if required)

#### 4. TERM OF PLAN

This Interim Recovery Plan will operate from August 2005 to July 2010 but will remain in force until withdrawn or replaced. If the taxon is still ranked as Endangered after five years, this IRP will be reviewed and if necessary, further recovery actions put in place.

#### 5. REFERENCES

- Brown, A., Thomson-Dans, C. and Marchant, N. (Eds) (1998) Western Australia's Threatened Flora. Department of Conservation and Land Management.
- CALM (2004) Towards a biodiversity conservation strategy for Western Australia : discussion paper. Department of Conservation and Land Management, Perth, Western Australia.
- Gill, A.M. and Nichols, A.O. (1989) Monitoring fire prone flora in reserves for nature conservation. In: "Fire Management on Nature Conservation Lands". Occasional Paper 1/89. Department of Conservation and Land Management, Perth, Western Australia.
- Hobbs, R.J. and Yates, C.J. (2003) Impacts of ecosystem fragmentation on plant populations: generalising the idiosyncratic. *Australian Journal of Botany* **51**: 471-488.
- Maslin, B.R. (2001) WATTLE [electronic resource]: acacias of Australia. Australian Biological Resources Study, Canberra.
- Nichols, O. (1998) Impacts of dieback-induced vegetation changes on native faunal communities in south-western Australia. Scope Item 7 In: Control of *Phytophthora cinnamomi* and Diplodina Canker in Western Australia. Final Report to the Threatened Species and Communities Unit, Biodiversity Group, Environment Australia, Department of Conservation and Land Management, Western Australia.
- Peters, R.L. and Darling, J.D.S. (1985) The greenhouse effect and nature reserves. *Bioscience* **35**(11): 707-717.
- Shearer, B. and Tippett, J. (1989) Jarrah dieback, the dynamics and management of *Phytophthora cinnamomi* in the jarrah (*Eucalyptus marginata*) forest of south-western Australia. Research Bulletin No.3. Department of Conservation and Land Management, Western Australia.
- Schwarten, T. (1995) The Biology and Ecology of Threatened *Daviesia* Species in Western Australia - Final Report, Kings Park and Botanic Garden, Western Australia.
- Stone, G.N., Raine, N.E., Prescott, M. and Willmer, G. (2003) Pollination ecology of acacias (Fabaceae, Mimosoideae). *Australian Systematic Botany* **16**: 103-118.
- Yates, C.J. and Broadhurst, L.M. (2002) Assessing limitations on population growth in two critically endangered *Acacia* taxa. *Biological Conservation* **108**(1): 13-26
- Weste, G. and Marks, G. (1987) The biology of *Phytophthora cinnamomi* in Australasian forests. *Annual Review of Phytopathology* **24**: 207-229.
- Wills, R. (1993) The ecological impact of *Phytophthora cinnamomi* in the Stirling Range National Park, Western Australia. *Australian Journal of Ecology* **18**: 145-159.
- Wilson, B., Newell, G., Laidlaw, W. and Friend, G. (1994) Impact of plant diseases on faunal communities. *Journal of the Royal Society of Western Australia* **77**: 139-144.
- World Conservation Union (2000) *IUCN red list categories prepared by the IUCN Species Survival Commission, as approved by the 51<sup>st</sup> meeting of the IUCN Council*. Gland, Switzerland.

## 6. TAXONOMIC DESCRIPTION

Spreading, open subshrub 0.2–0.4 m high. Stems slender, dark grey. Branchlets densely pubescent. Stipules setaceous, 5–7 mm long, recurved. Phyllodes crowded, linear, 11–17 mm long, 1–1.5 mm wide, narrowed at base, excentrically rostellate, dark green, glabrous but pulvinus pubescent adaxially; midrib near abaxial margin and prominently raised, the 2-nerved adaxial margin thick and nerve-like. Inflorescences rudimentary, 1-headed racemes with axes <0.5 mm long; peduncles 8–13 mm long, glabrous, recurved in fruit; basal bract cucullate-navicular; heads globular, 2.5–3 mm diam., 12–16-flowered, light golden. Flowers 5-merous; sepals free. Pods subterete, 10–15 mm long, thinly crustaceous, blackish. Seeds longitudinal, oblong-elliptic to ovate, 2–2.5 mm long, c. 1.5 mm wide, shiny, dark brown; aril pileiform.

